

## Physical and biological triggers of ventilator-induced lung injury and its prevention

L. Gattinoni, E. Carlesso, P. Cadringer, F. Valenza, F. Vagginelli, D. Chiumello

*Physical and biological triggers of ventilator-induced lung injury and its prevention.*  
L. Gattinoni, E. Carlesso, P. Cadringer, F. Valenza, F. Vagginelli, D. Chiumello.  
©ERS Journals Ltd 2003.

**ABSTRACT:** Ventilator-induced lung injury is a side-effect of mechanical ventilation. Its prevention or attenuation implies knowledge of the sequence of events that lead from mechanical stress to lung inflammation and stress at rupture.

A literature review was undertaken which focused on the link between the mechanical forces in the diseased lung and the resulting inflammation/rupture.

The distending force of the lung is the transpulmonary pressure. This applied force, in a homogeneous lung, is shared equally by each fibre of the lung's fibrous skeleton. In a nonhomogeneous lung, the collapsed or consolidated regions do not strain, whereas the neighbouring fibres experience excessive strain. Indeed, if the global applied force is excessive, or the fibres near the diseased regions experience excessive stress/strain, biological activation and/or mechanical rupture are observed. Excessive strain activates macrophages and epithelial cells to produce interleukin-8. This cytokine recruits neutrophils, with consequent full-blown inflammation.

In order to prevent initiation of ventilator-induced lung injury, transpulmonary pressure must be kept within the physiological range. The prone position may attenuate ventilator-induced lung injury by increasing the homogeneity of transpulmonary pressure distribution. Positive end-expiratory pressure may prevent ventilator-induced lung injury by keeping open the lung, thus reducing the regional stress/strain maldistribution. If the transpulmonary pressure rather than the tidal volume per kilogram of body weight is taken into account, the contradictory results of the randomised trials dealing with different strategies of mechanical ventilation may be better understood.

*Eur Respir J 2003; 22: Suppl. 47, 15s–25s.*

Institute of Anaesthesia and Intensive Care,  
University of Milan, Policlinico Hospital,  
IRCCS, Milan, Italy.

Correspondence: L. Gattinoni  
Istituto di Anestesia e Rianimazione  
Ospedale Maggiore Policlinico-IRCCS  
Via Francesco Sforza, 35  
20122 Milano  
Italy  
Fax: 39 0255033230  
E-mail: gattinon@policlinico.mi.it

Keywords: Acute respiratory distress syndrome  
elastance  
respiratory mechanics  
stress/strain  
ventilator-induced lung injury

Received: February 25 2003  
Accepted after revision: July 14 2003

In the 1970s, the typical setting for mechanical ventilation in acute respiratory distress syndrome (ARDS) was a tidal volume ( $V_T$ ) of 12–15 mL·kg body weight<sup>-1</sup> with a positive end-expiratory pressure (PEEP) of 5–10 cmH<sub>2</sub>O.

"We ventilated thousands of patients in this way and the only side effect was hypocapnia" was a statement made by PONTOPPIDAN *et al.* [1] in the *The New England Journal of Medicine* in 1972. However, since the mid-1970s, there has been progressive recognition of the potential harm of mechanical ventilation. The recognition that the ventilatable lung in ARDS is small, and not stiff (the baby lung concept [2]), made obvious the possible mechanical harm of using a high  $V_T$  in a little lung, and led to concepts such as permissive hypercapnia [3], *i.e.* to more gentle lung treatment. In parallel, it was recognised *ex vivo* [4] and *in vivo* [5] that, irrespective of mechanical rupture, mechanical ventilation may induce a complex biological response, with release of inflammatory and anti-inflammatory mediators. This led to the concept of biotrauma, which is widely, although not universally [6], accepted. In the present article, the authors present their understanding of the initial physical and biological events which trigger ventilator-induced lung injury (VILI) and its clinical consequences, without intending to either provide an exhaustive review of full-blown VILI or preclude or invalidate other interpretations.

### The distending force of the lung and its distribution in healthy and diseased lung parenchyma

The distending force of the lung per unit area, *i.e.* the pressure, is that applied to the visceral pleura. This is the transpulmonary pressure ( $P_L$ ), which is the difference between the pressure inside the alveoli and the pleural pressure ( $P_{pl}$ ). Unfortunately, in normal clinical practice, it is usual to consider the plateau or airway pressure ( $P_{aw}$ ) as the distending force of the lung. Under static conditions,  $P_{aw}$  closely reflects the intra-alveolar pressure, which, in part, is spent to inflate the lung, and, in part, to inflate the chest wall. In a simple model that ignores blood shift, lung inflation is considered to be equivalent to chest wall inflation. The relationship between  $P_{aw}$  and  $P_{pl}$ , which determines the  $P_L$ , depends on the relative mechanical characteristics of lung and chest wall, which is convenient to express as elastance (pressure/volume).

The elastance of the respiratory system ( $E_{rs}$ ) is the  $P_{aw}$  required to inflate the respiratory system to 1 L above its resting position under static conditions. Indeed,  $P_{aw}$  equals the sum of the pressure used to inflate the lung ( $P_L$ ) and the one used to inflate the chest wall ( $P_{pl}$ ):

$$P_{aw} = P_L + P_{pl}, \quad (1)$$

and

$$E_{rs} = E_L + E_w, \quad (2)$$

where  $E_L$  and  $E_w$  are the elastances of the lung and chest wall. Accordingly,

$$P_L = P_{aw} E_L / E_{rs}. \quad (3)$$

Under normal static conditions, since  $E_L$  is approximately the same as  $E_w$ , whatever pressure is applied to the alveoli, it results in a change in  $PL$  of  $\sim 50\%$  ( $E_L/E_{rs}=0.5$ ).

For many years, unfortunately, any change in  $E_{rs}$  was attributed to  $E_L$ . However, there is now consistent evidence that  $E_w$  may be altered in ARDS patients [7–10] due either to individual patients' anatomical characteristics (body size and weight) or the nature of the disease leading to ARDS. It has been shown that abdominal pressure, which directly increases  $E_w$ , is frequently altered in abdominal diseases associated with ARDS, such as peritonitis or bowel ischaemia [11, 12]. Indeed the  $E_L/E_{rs}$  ratio, which determines the relationship between  $P_{aw}$  and the resulting  $PL$  (Equation 3), may range in acute lung injury/ARDS from 0.3 to 0.8 or more, greatly different from the normal value of 0.5. Since VILI occurs due to distension of the lung, which depends on  $PL$ , it follows that its measurement or estimation is a key issue, as, for the same applied  $P_{aw}$ ,  $PL$  and its potential harm to the lung may vary greatly between patients.

### Pressure transmission throughout the lung

The  $PL$  is applied to the visceral pleura and has to be transmitted throughout all lung regions. The force-transmitting system is the lung's fibrous skeleton, which consists of two components, the axial fibres, anchored to the hilum, which run along the branching airways down to the level of alveolar ducts, and the peripheral fibre system, anchored to the visceral lung pleura, which penetrates centripetally into the lung down to the acini. The two systems are linked at the alveolar level by the alveolar septal fibres [13]. The lung's fibrous skeleton consists mainly of elastin and collagen fibres, which are intimately associated with each other. Other load-bearing force elements connecting to the pulmonary interstitium are actin and myosin microfilaments originating from myofibroblasts.

### Pressure, stress/tension and strain

The strict definition of stress and strain is beyond the purpose of the present article and may be found elsewhere [14]. However, in its simplest definition, in a monodimensional structure (such as a string/spring pair), stress (or tension) may be defined as the force per unit area which develops in a structure as a reaction to an applied external force of the same entity but opposite direction, *i.e.*  $\sigma n = \Delta F / \Delta S$ , where  $\sigma n$  is stress,  $\Delta F$  change in force and  $\Delta S$  the reference surface to which the force is applied. This internal force, cutting the structure by an ideal surface, may be resolved into three components, one of which is normal to the plane (normal stress) and two of which are tangential (shear stresses). The deformation of the structure (if any) due to the applied force is called strain. In the simplest monodimensional structure under traction, strain is defined as the ratio of the change in length of the structure ( $\Delta L$ ) to its length in the resting position ( $L_0$ ), *i.e.*  $\epsilon = \Delta L / L_0$ , where  $\epsilon$  is strain. Indeed, stress and strain are the natural response of a structure to an applied force. In the lung, during mechanical ventilation, stress and strain are periodically changing variables characterised

by maximal and minimal values (end-inspiratory and end-expiratory  $PL$  for stress, and end-inspiratory and end-expiratory lung volume (EELV) for strain) at a given frequency and amplitude (difference between maximal and minimal value).

When  $PL$  is applied to the visceral pleura surface by the inspiratory muscles or ventilator, this applied force, under static conditions, is equal to the sum of the forces developed within the lung parenchyma. Part of these forces are borne by the alveolar air/liquid interface, which, in the presence of surfactant, are very low up to 80% of total lung capacity [15], whereas the remaining forces are shared by the fibre system of the lung's fibrous skeleton. With lack of surfactant, the liquid/air interface bears more force and conceptually works as an "added force-bearing fibre system". Indeed, each fibre experiences a stress/tension according to the force it has to bear. In a homogeneous lung, every fibre shares an equal proportion of the total force applied and develops an equal tension and equal strain. If some of the fibres are destroyed (as in emphysema), fewer fibres have to share the applied force and experience greater stress and strain. However, if part of the parenchyma remains collapsed during inflation or cannot expand, as in consolidated pneumonia, the interwoven fibres in the diseased region bear the force and are in tension but do not strain. The fibres connected to the not expandable region, however, have to carry a greater force load, with greater tension and distortion (fig. 1). These concepts have been developed, on a theoretical background, by MEAD *et al.* [16], who, in a simplified model in which the volume of the collapsed region was one tenth of the volume that this region would have occupied if not collapsed, computed that, for an applied pressure of 30 cmH<sub>2</sub>O, the resulting tension in the neighbouring region would have been 140 cmH<sub>2</sub>O. Independently of the precision of this computation, it is clear that, when the stress and strain are not homogeneously distributed, greater tension and distortion develop in some regions and the order of magnitude of fibre tension may be such to lead to stress at mechanical rupture.

### Targets of injury

As discussed above, it appears that the triggers of VILI are the mechanical forces which globally (excessive applied  $PL$ ) or locally (stress/strain maldistribution due to lung inhomogeneity) cause mechanical alterations of the lung parenchyma which range from excessive and nonphysiological strain up to stress at rupture. Indeed, the three main targets of injury are the fibre systems in the extracellular matrix, alveolar cells and lung capillaries. The small airways may also be affected [17, 18], but have scarcely been studied and so are not discussed further.

### Fibre system

As the most important components of the lung's fibrous skeleton are the collagen and elastin fibres, it is appropriate to briefly summarise their mechanical characteristics (fig. 2, table 1). The simplest approach to elastin/collagen interaction is to consider the elastin a spring in parallel with a folded string of collagen. The elastic behaviour of the simple unit is shown in figure 2a. When an external force is applied, the spring (elastin) is the force-bearing element and develops stress and strain according to its mechanical characteristics. The string (collagen) develops its stress when completely unfolded. Since the collagen is almost inelastic, it works as a stop-length fibre, preventing further strain of the elastin/collagen unit. However, different units may have different elastic constants




Fig. 1. – Model of fibrous network showing three fibre lines in parallel (A–C), with each line containing three springs (1–3). a) Resting position. b) When a force ( $F$ ) is applied and the system is homogeneous, each line bears a third of the force ( $0.33F$ ) and each spring in each line carries the same force ( $0.33F$ ). Thus the strain is equally distributed within the lines and the springs. c) If one element is consolidated (inclusion), it carries the force but does not strain. The connected springs experience greater strain. d) In the case of rupture (of line B), the line does not carry any stress/strain; however, the lines in parallel undergo greater stress and strain. In mathematical terms, each spring has stiffness  $K$ , except spring B2, whose stiffness is  $MK$  and which determines the different behaviours of the model. When  $F$  is applied, the model behaves according to the following equations:  $\Delta L = F(6M+3)/(K(7M+2))$ ;  $\Delta L_{A1} = \Delta L_{A2} = \Delta L_{A3} = \Delta L_{C1} = \Delta L_{C2} = \Delta L_{C3} = \Delta L/3$ ;  $\Delta L_{B1} = \Delta L_{B3} = M\Delta L/(2M+1)$ ;  $\Delta L_{B2} = \Delta L/(2M+1)$ ;  $F_A = F_C = F(6M+3)/(21M+6)$ ; and  $F_B = 3MF/(7M+2)$ . If  $M=1$  the system is homogeneous; if  $M=0$ , line B is broken; and, finally, if  $M \rightarrow \infty$ , B2 behaves as an inclusion.  $\Delta L$ : change in length of structure;  $L_0$ : total length in resting position.

and different stop lengths. When connecting different units with different characteristics in series, the overall behaviour results from the contribution of the simple elements (fig. 2b and c). The situation is more complex when considering the network of spring/string units connected in series and parallel. Assuming a statistical density distribution of elastin/collagen mechanical characteristics, models have been developed, which describe the mechanical behaviour of the system as tested *ex vivo* (fig. 2d) [19, 20]. Considering the mechanical behaviour of the whole fibre system and its tissue/air ratio, it appears that the order of magnitude of the stress at rupture is  $\sim 100 \text{ cmH}_2\text{O}$ .

Alveolar cells

Three-quarters of all lung cells (by volume) are located in gas exchange regions. Although type II epithelial cells are located in the alveolar corners, type I epithelial cells (about  $\sim 90\%$  of the alveolar surface) are flat and wide, and the same cell may encompass, in a sandwich-like fashion, approximately four endothelial cells. It is worth emphasising that, in most of the alveolar structure, type I epithelial cells share a common basal membrane with endothelial cells, suggesting mechanical coupling. The fibre system and associated fibroblasts as well as filaments of actin and myosin, which all contribute to mechanical support [21], are located in the basal membrane (extracellular matrix) to which both the epithelial and endothelial cells are anchored *via* integrins.

When a distending force is applied to the fibre system, which primarily bears the load, all of the anchored cells have to accommodate their shape to the new surface. Obviously, there is a continuum from physiological deformation up to plasma cell break (stress failure). The interaction between the mechanical deformation and the biological reaction has been extensively investigated in cell culture (excellently reviewed in [22–28]). However, it is unlikely that the stress/strain relationship in cell culture is equivalent to that *in vivo* due to the complex architecture of the alveolar wall as well as phenomena such as alveolar cell unfolding at high volume [29]. Therefore, it is hard to translate lung volume changes into cell strain changes. However, despite these limitations, the present authors believe that the available data, obtained in cultures of different alveolar cells, may be considered as offering a unique perspective, which is internally consistent.

As summarised in figure 3, the cells react to deformation, first reinforcing the plasma membrane by recruiting intracellular lipids to the cell surface, a phenomenon called deformation-induced lipid trafficking [30, 31]. In the meantime, the "mechanosensors", *i.e.* the integrins, the cytoskeleton and ion channels, transduce the mechanical signal in biochemical events, *via* a complicated network of signalling molecules [26–28]. The final result of an "intermediate" nonphysiological strain, not such as to produce physical rupture of the alveolar wall, is the reinforcement/sealing of the plasma cell membrane [32], as well as, *via* mechanosensors, upregulation of inflammatory cytokines and, possibly,  $\text{Ca}^{2+}$ -mediated cell contraction [33].

The results of various experiments in different cell cultures are summarised below. At a strain causing a 12% increase in surface area, it has been shown that human macrophages, *via* nuclear factor- $\kappa\text{B}$ , produce interleukin (IL)-8 [34], a cytokine of the CXC chemokine family, which is the most powerful chemoattractant for neutrophils. At the same level of strain, macrophages produce metalloproteins, which remodel the extracellular matrix. Interestingly, no alveolar cells tested at this level of strain produced any other cytokine, including tumour necrosis factor- $\alpha$ . At 17–18% linear strain change




Fig. 2. – Mechanical characteristics of elastin and collagen: Young module (stress/strain) (see table 1). Stress/strain relationship of: a) one; b) two; and c) seven elastin/collagen pairs of elasticity  $K$  (—) and maximal length  $L$  (.....). The elastin/collagen pairs have the same  $K$  but different  $L$  and are connected in series (○: stress/strain at which the various collagen fibres become unfolded and reach their stop length). d) Stress/strain relationship obtained in tissue strips (—: behaviour of whole system; - - - -: elastic behaviour of elastin fibres; - - - -: elastic behaviour of collagen fibres; .....: elastin stress at rupture limit). It appears that the stress at rupture (indicated by vertical arrow) is 1,000 kPa. As the ratio between tissue and air is  $\sim 1:100$ , the stress at rupture in the lung should be  $\sim 1,000/100$  kPa, *i.e.* 100 cmH<sub>2</sub>O. AU: arbitrary units. (d) adapted from [19].

Table 1. – Mechanical properties of collagen and elastin fibres

	Young module MPa	Stress at rupture MPa	Strain at rupture %	Elastic limit %	Content %
Collagen	1000	50–100	10	1–2	2
Elastin	0.6	1	100	60	10

(which should correspond to  $\sim 37\%$  surface change), it has been shown that human endothelial cells produce metallo-proteins [35]. At 30 and 40% strain, A549 epithelial cells produce IL-8 [36, 37], whereas, at 50% surface strain, which should correspond, in that experimental set-up with rat epithelial cells, to a volume change greater than total lung

capacity *in vivo*, 70% cell death has been reported [38]. These data refer to the magnitude of strain (a rough equivalent of end-inspiratory lung volume). However, it has been shown, both *in vitro* and *in vivo*, that the duration of strain, as well as its amplitude and frequency, may increase the injury [39–42]. Interestingly, for the same magnitude of strain, reducing its




Fig. 3.—Simplified scheme showing cell reaction to mechanical stress (-----: separates different stages in reaction). DILT: deformation-induced lipid trafficking; MAPK: mitogen-activated protein kinase; NF-κB: nuclear factor-κB; IL: interleukin. (Data from [25, 28<sup>#</sup>, 30, 31<sup>\*</sup>]).

amplitude (by increasing baseline strain) reduces epithelial cell injury [40]. Indeed, from the bulk of data, it appears that cyclic nonphysiological strain on alveolar cells induces release of IL-8 and metalloproteins, whereas cell death occurs when the strain exceeds the total volume capacity. It is tempting to speculate that the possible first trigger of the biological reaction is IL-8, the most powerful chemokine for neutrophil activation.

The role of IL-8 as a first cytokine responsible for the further sequence of events that leads to inflammation, through neutrophil recruitment, has been recently underlined in a mouse model. BELPERIO *et al.* [43] showed, in mice ventilated at a *V*<sub>T</sub> of 6 (low-strain group) and 12 (high-strain group) mL·kg body weight<sup>-1</sup>, that neutrophil activation was increased compared to the spontaneously breathing control group and that neutrophil activation was proportional to the strain. The strain/injury was associated with an increase in levels of CXC2 chemokine (a murine equivalent of IL-8) and its receptors. Blocking CXC2 or its receptors with specific antibodies, or using knockout mice for CXC2 receptors, did not induce neutrophil activation and attenuated VILI. The bulk of data strongly support a cause/effect relationship between strain, IL-8 production, neutrophil activation and VILI.

*Lung capillaries*

The fibre system provides mechanical support for the pulmonary blood vessels. In the alveolar septum, where

the axial and peripheral fibre systems are interconnected, the alveolar capillary network is interwoven with the meshwork of septal fibres. It has been known for a long time that excessive strain of the lung structures, for minutes or days, depending on animal species, causes pulmonary oedema of varying degree (the closed chest approach results in less oedema than the open chest approach when applying the same *P*<sub>aw</sub>), associated with gas exchange impairment, hyaline membrane formation and neutrophil infiltration [44–49]. The key issue is understanding how the excessive mechanical strain causes oedema. When the alveolar fibre network experiences a nonphysiological excessive strain, the capillary meshwork of the alveolar septum flattens, whereas the corner vessels maintain or increase their patency. The final result is increased resistance to blood flow, which leads to increased pulmonary artery pressure. This, in turn, causes an increased filtration rate in excess of the increased lymph flow, with fluid accumulation in the interstitial spaces. Indeed, part of the oedema due to high pressure/volume ventilation is "hydrostatic" in nature [50].

However, the excessive strain also causes increased permeability of the capillary network [45, 46]. This was initially attributed to a "stretched pore phenomenon", a passive process due to the increased hydrostatic pressure forcing the loose connections between the endothelial cells. However, it has also been shown that intercellular gaps may occur at high transmural capillary pressure despite the intact extracellular matrix [51–53]. This active process, probably involving cell contraction [54], is possibly due to Ca<sup>2+</sup> influx through

mechanically gated calcium channels. The increase in intracellular  $\text{Ca}^{2+}$  level has multiple effects that may influence permeability, including increased actin/myosin filament tension. Blocking the  $\text{Ca}^{2+}$  influx using gadolinium, an inhibitor of stretch-activated cation channels [55], or preventing actin/myosin filament contraction [56] significantly reduces endothelial permeability. Indeed, opening of intracellular/intercellular gaps, remodelling of the cytoskeleton and active cell contraction may all contribute to increased permeability. Moreover, full-blown inflammation, in which neutrophils are recruited, may very well induce, through a variety of mediators, increased endothelial permeability [54]. At intermediate degrees of strain, such mechanosignalling may be the trigger of VILI. When the applied mechanical stress is very high, however, the extracellular matrix may break, the inflammatory process being a consequence rather than the initiator/associated trigger of the observed damage. Indeed, it is quite clear that, depending on the applied stress/strain, pulmonary oedema may occur with or without inflammation.

It is worth underlining, however, the harmful interaction between excessive alveolar strain, pulmonary artery pressure and lung capillary blood volume. First, high capillary pressure may induce stress failure with increased permeability [53] and any increase in pressure increases oedema formation [57]. Moreover, it has been shown that cyclic changes in perivascular pressure surrounding extra-alveolar vessels due to mechanical ventilation cause greater oedema than isolated phasic elevation of pulmonary artery pressure without mechanical ventilation [41]. Finally, not only elevated but also low capillary pressure may damage the lung. Indeed, low capillary pressure may facilitate collapse and decollapse of the alveolar capillaries, with possible stress failure, whereas, in the meantime, it may increase transmural pressure in extra-alveolar capillaries, with increased oedema formation [58].

### Ventilator-induced lung injury and species

As most of the studies on VILI have been performed in animals, it is worth discussing some of the important differences between the various animal species. In some experiments, VILI was induced by using different  $V_T$  normalised to body weight (in kilograms) (table 2) [43, 48, 49, 59, 60]. Unfortunately, lung volume, alveolar size and body weight are not linearly related across the various species. In figure 4, based on anatomical data reported by MERCER *et al.* [64], alveolar diameter changes, which reflect the tension to which the fibres of the lung's fibrous skeleton are subjected, are shown relative to  $V_T$  per kilogram of body weight. As shown, a  $V_T$  of  $10 \text{ mL}\cdot\text{kg}^{-1}$  in normal humans induces an increase in alveolar diameter of  $\sim 10\%$ . In mice, the same normalised  $V_T$  induces an alveolar diameter increase of  $\sim 40\%$ , which, in humans, would correspond to a  $V_T$  of  $\sim 45 \text{ mL}\cdot\text{kg}^{-1}$ . In other experimental studies on

VILI, the injurious strategy was applied using high  $P_{aw}$  rather than high  $V_T$ . However, the relationship between  $P_{aw}$  and  $PL$ , the real trigger of VILI, depends on the ratio between  $EL$  and  $E_{rs}$  (Equation 3). This ratio ( $EL/E_{rs}$ ) varies widely across species, from near 1 in mice to  $\sim 0.5$  in normal humans (table 2).

If  $PL$  is taken into account, the distinction between volutrauma and barotrauma vanishes. In the experiments of DREYFUSS *et al.* [65], which led to the volutrauma concept, the high  $P_{aw}/\text{low } V_T$  was obtained by increasing the  $E_w$  by strapping the rat thorax with rubber bands. In this case, the  $PL$ , despite the high  $P_{aw}$ , was greatly decreased, and, obviously, the lung damage was lesser than in rats ventilated at the same  $P_{aw}$  with normal  $E_w$ .

Indeed, from the bulk of the data, it appears that, to produce VILI in normal lung, high  $PL$  need to be used, irrespective of whether volume control or pressure control ventilation are used.

However, in clinical practice, the great concern is VILI in an already diseased lung. Interestingly, the few experimental data available suggest that VILI may be induced in diseased lung at lower  $V_T/P_{aw}$  than in normal lung [66, 67]. This is quite understandable if the relationship between  $E_{rs}$  and  $EELV$  is taken into account. In ARDS, the lung is "small" rather than "stiff" [2], and  $E_{rs}$  is a function of ventilatable  $EELV$  [68], specific  $E_{rs}$  ( $E_{rs}/EELV$ ) being near normal [69]. Indeed, the tension throughout the lung parenchyma depends on the ratio between the  $V_T$  and  $EELV$  to which the  $V_T$  is delivered. For example, in a human with severe ARDS with an  $EELV$  of 500 mL (the baby lung [2]), a  $V_T$  of 500 mL induces approximately the same tension as a  $V_T$  of 2,500 mL in a human with a normal  $EELV$  of 2,500 mL.

Indeed, whereas  $PL$  may be considered the rough clinical equivalent of stress, the  $V_T/EELV$  ratio may be viewed as the rough clinical equivalent of strain. Stress ( $PL$ ) and strain ( $V_T/EELV$ ) are linked by specific  $EL$  ( $EL_{sp}$ ) according to the formula

$$EL_{sp} = (\Delta PL / \Delta V_T) EELV. \quad (4)$$

It, therefore, follows that

$$\Delta PL = EL_{sp} \Delta V_T / EELV. \quad (5)$$

This indicates that considering the  $PL$  or the  $V_T/EELV$  ratio (*i.e.* the ratio between the inflation of the whole lung, or any given region, and its resting volume) are two ways of looking at the same reality, thus reunifying the concepts of barotrauma (not  $P_{aw}$  alone but  $PL$ ) and volutrauma (not  $V_T$  alone but  $V_T/EELV$ ), according to basic physiology. It may also be useful to point out that  $EL_{sp}$  is a kind of "Young module" for the lung (*i.e.* stress/strain) and stress is the independent variable in pressure control ventilation, whereas strain is the independent variable in volume control ventilation.

Table 2. – Experimental ventilator-induced lung injury in normal lungs

First author [Ref.]	Species	$P_{aw}$ cmH <sub>2</sub> O	$V_T$ mL·kg body weight <sup>-1</sup>	Alveolar diameter deformation %	$E_{rs}^{\#}$ cmH <sub>2</sub> O·mL <sup>-1</sup>	$E_w$ cmH <sub>2</sub> O·mL <sup>-1</sup>	$EL$ cmH <sub>2</sub> O·mL <sup>-1</sup>
KOLOBOW [48]	Sheep	50	50–70	–	0.050 [61]	0.020	0.030
BROCCARD [59]	Dog	44	77	–	0.027	0.004	0.023
NISHIMURA [60]	Rabbit	24 <sup>†</sup>	31	>100	0.230 [18]	0.030	0.200
WEBB [49]	Rat	40	40	70	3.400 [62]	0.400	3.000
BELPERIO [43]	Mouse	40	24	72	13.70 [63]	Near 0.00	13.70

$P_{aw}$ : airway pressure;  $V_T$ : tidal volume;  $E_{rs}$ : elastance of the respiratory system;  $E_w$ : elastance of the chest wall;  $EL$ : elastance of the lung. <sup>#</sup>: additional references provided as data not given in articles cited; <sup>†</sup>: transpulmonary pressure.






Fig. 4.—Relative increase in alveolar diameter as a function of tidal volume ( $V_T$ ) normalised to body weight in different species (—: human (1.00); - - - -: monkey (1.40); - - - -: baboon (1.58); - - - -: rat (2.13); - - - -: hamster (2.23); - - - -: rabbit (3.08); - - - -: mouse (3.18)). Alveolar diameter changes were computed on the assumption that the relationship existed in a sphere, *i.e.*  $\% \Delta \text{diameter} = 1 + (1 + \% \Delta \text{volume})^{0.33}$ . The strain index, *i.e.* the alveolar diameter change relative to that expected in humans, at a  $V_T$  of 10 mL·kg body weight<sup>-1</sup> (.....) is indicated above in parenthesis for each species. (Baseline data were derived from [64]).

**Prevention of ventilator-induced lung injury**

VILI is the result of a sequence of events, which begins with mechanical alteration of the lung parenchyma, due to excessive global and/or regional stress/strain. If the resulting tension in the structure reaches the limits of stress at rupture, the structures are destroyed (alveolar wall and capillaries). If the tension is lower than these limits, but nonphysiological, a biological reaction occurs, probably involving, first, the macrophages, with IL-8 production, from which the inflammatory cascade, through neutrophil recruitment, fully develops. Patients with VILI present with typical lung inflammation, with all its biochemical, histological and pathological characteristics. Indeed, VILI may be prevented or attenuated by interfering with the sequence of biological reactions leading to the inflammatory response (chemokine antibodies [43], steroids [70], *etc.*). However, since nonphysiological stress and strain appear to be the first trigger of VILI, the possibilities available for limiting/preventing the excessive regional and global stress and strain of the ARDS lung, *i.e.* prone position, PEEP and low  $V_T$ , are discussed.

*Prone positioning*

Thus far,  $PL$  has been considered as being uniformly distributed; however, it is well known that, in both humans and experimental animals, there is a gradient of  $PL$  along the vertical axis, with the nondependent lung regions experiencing greater  $PL$  and greater tension of the fibres of the lung's fibrous skeleton than dependent lung regions. This phenomenon is enhanced in a nonhomogeneous lung. It was found, for example, that, in an oleic acid model in the supine position, the difference in  $PL$  between nondependent and dependent regions was as great as 10 cmH<sub>2</sub>O [71]. In the prone position, in both humans [72] and experimental settings [73], regional inflation is more uniformly distributed along the vertical axis, indicating a significant reduction in the  $PL$  gradient. This suggests that the stress and strain are more




Fig. 5.—Relationships between tidal volume ( $V_T$ ), plateau pressure ( $P_{aw}$ ; - - - -) and transpulmonary pressure ( $P_L$ ; —) in a model of a 70-kg human at 10 cmH<sub>2</sub>O positive end-expiratory pressure with constant compliance of the respiratory system of: a) 50; b) 30; and c) 20 mL·cmH<sub>2</sub>O<sup>-1</sup>, *i.e.* elastances of the respiratory system ( $E_{rs}$ ) of 0.02, 0.033 and 0.05 cmH<sub>2</sub>O·mL<sup>-1</sup> respectively.  $P_{aw}$  were computed using  $P_{aw} = V_T E_{rs}$ . At each  $P_{aw}$ , the  $P_L$  resulting from elastances of the lung ( $E_L$ )/ $E_{rs}$  ranging 0.3–0.8 (■: 0.3; □: 0.4; ●: 0.5; ○: 0.6; ▲: 0.7; △: 0.8), *i.e.*  $P_L = P_{aw} E_L / E_{rs}$ , were computed. The dangerous  $P_L$  was arbitrarily set at 15 cmH<sub>2</sub>O (.....), which correspond to ~70–75% of total lung capacity in normal humans; the area below this represents the "safe zone". Depending on  $E_{rs}$  and  $E_L/E_{rs}$ , a  $V_T$  of 6 or 12 mL·kg body weight<sup>-1</sup> could be either harmful or safe.

homogeneously distributed within the lung parenchyma, and this is the rational basis for the possible effectiveness of prone positioning in attenuating VILI, as shown experimentally in dogs [59] and rabbits [60]. Unfortunately proof is still lacking in patients that prone positioning affects outcome. However, in a subgroup of acute lung injury/ARDS patients treated with high-volume ventilation ( $V_T$  of  $\geq 12$  mL·kg body weight<sup>-1</sup>), the mortality rate of patients ventilated in the supine position was significantly greater (almost double) than that observed in patients ventilated while prone [74].




Fig. 6.—Distribution of: a, c, e, g) tidal volume ( $V_T$ ); and b, d, f, h, i) plateau pressure ( $P_{aw}$ ) (—: treated group; .....: controls) in the available randomised trials (computed from the reported mean $\pm$ SD assuming a Gaussian distribution): a, b) 432 treated/429 controls [81]; c, d) 58 treated/58 controls [79]; e, f) 60 treated/60 controls [77]; g, h) 26 treated/26 controls [80]; and i) 29 treated/24 controls [78].

### Positive end-expiratory pressure

As PEEP unavoidably results in an increase in mean  $PL$ , leading to greater stress of the lung parenchyma, it is quite surprising, at first sight, that it is so effective in many (but not all) circumstances in attenuating VILI, as reviewed extensively by DREYFUSS and SAUMON [75].

A possible explanation is that, if a lung region is collapsed/consolidated and does not expand during inspiration, the fibres of the neighbouring open regions show increased tension and strain. Indeed, if PEEP is effective in keeping open the collapsed region, the applied force is shared by more fibres with more uniform distribution of stress and strain. Interestingly, the positive effect of PEEP in protecting against VILI have been described in animal models with great potential for lung recruitment, in which PEEP is effective in keeping open most of the lung, thus attenuating the stress/strain maldistribution. It may be wondered, however, whether VILI can be prevented by PEEP when most of the lung is consolidated and the potential for lung recruitment is very low, as in diffuse pneumonia [11]. Although not proved, it is possible that, in this setting, PEEP simply increases

total stress without affecting stress/strain maldistribution. Interestingly, there are no reports showing positive effects of PEEP on outcome in patients, and the PEEP/outcome relationship is still a controversial issue. However, this is not surprising since PEEP was tested in patients with varying probable potentials for lung recruitment. It is possible that the positive effect of PEEP in the patient subgroup with a high potential for lung recruitment was obscured by its negative or zero effects in the subgroup of patients with a low potential for lung recruitment.

### Low tidal volume

Following consensus conference suggestions [76], since the late 1990s, several studies [77–81] have been performed in order to investigate the effects on outcome of low *versus* high  $V_T$ . These studies, although based on the same rationale (gentle lung treatment), were of different power and design. Three studies compared  $V_T$  of 7 and 10–10.5 mL·kg ideal body weight<sup>-1</sup> and were not able to show any difference in outcome [77, 79, 80]. The study of AMATO *et al.* [78] compared



two ventilatory strategies, high PEEP/low  $\dot{V}_T$  and low PEEP/high  $\dot{V}_T$ . This study showed an impressive difference in mortality between the two strategies but was criticised, mainly because of the high (70%) mortality in the high  $\dot{V}_T$ /low PEEP group. The last study of the series was performed by the National Institutes of Health (NIH) network and tested, in an adequately powered trial, the mortality differences between patients treated with  $\dot{V}_T$  of 6 and 12 mL·kg body weight<sup>-1</sup> [81]. The results showed a significantly different outcome, with ~9% decrease in absolute mortality in the 6 mL·kg body weight<sup>-1</sup> group. In most of these studies, with the exception of the study of AMATO *et al.* [78], the "safety limit" for  $P_{aw}$  was set at 35 cmH<sub>2</sub>O. The contradictory results generated a lot of controversy, to the point that a recent meta-analysis claimed that the NIH network ventilation at a  $\dot{V}_T$  of 6 mL·kg body weight<sup>-1</sup> was unsafe, suggesting that the relationship between  $\dot{V}_T$  and outcome is U-shaped, with greater risk of mortality associated with low as well as high  $\dot{V}_T$  [82].

Before discussing the results of the available clinical studies, it is worth recognising that, if two different kinds of ventilation produce different outcomes, as in the NIH network ARDS trial, this probably means that the "amount of VILI" associated with the two different kind of ventilation is also different. However, since the real cause of VILI is the PL, it is evident that measuring  $\dot{V}_T$  is associated with a great deal of confounding variables. A given  $\dot{V}_T$  causes different  $P_{aw}$  depending on the  $E_{rs}$  ( $P_{aw}=E_{rs}\dot{V}_T$ ). A given  $P_{aw}$ , in turn, produces different PL according to the ratio of EL to  $E_{rs}$  ( $PL=P_{aw}EL/E_{rs}$ ). This indicates that the relationship between  $\dot{V}_T$  and the resulting PL may be highly variable. Indeed, it is quite obvious that, in a relatively small population, randomisation may be unable to distribute equally between groups of different EL and  $E_w$  present in the population under study. This bias should be attenuated in a large population. However, even in this case, the linkage between  $\dot{V}_T$  and PL is weak. This is underlined in figure 5, where PL is plotted as a function of  $\dot{V}_T$  normalised to body weight, for an ideal 70-kg human at 10 cmH<sub>2</sub>O PEEP, for a range of  $E_{rs}$  and  $EL/E_{rs}$ . Despite the obvious oversimplification of assuming, in this model, a linear volume/pressure curve, and setting the "dangerous PL" arbitrarily at 15 cmH<sub>2</sub>O (~70–75% of total lung capacity in normal humans), it is quite evident that potentially dangerous PL may be associated with a great variety of  $\dot{V}_T$  normalised to body weight as well as with  $P_{aw}$  well below the suggested "safety limit of 32 cmH<sub>2</sub>O" [83].

The  $\dot{V}_T$  distributions in the available randomised trials are depicted in figure 6. The distributions of  $P_{aw}$ , which is related to PL more than to  $\dot{V}_T$  (Equation 3), are also shown in figure 6. A huge overlap in the  $P_{aw}$  observed in the different studies is evident. However, the discussion between supporters of the importance of  $\dot{V}_T$  versus supporters of the importance of  $P_{aw}$  as VILI determinant does not seem to have a sound physiological basis as PL is the main determinant of VILI, and it has never been controlled in any randomised trial.

### Conclusions

On the basis of the available experimental and clinical data, the following conclusions may be drawn. Ventilator-induced lung injury is due to excessive global/regional stress/strain and affects the relatively healthier regions of the lung, since consolidated lung regions are not distended. The excessive stress/strain appears to activate first macrophages and subsequently neutrophils *via* interleukin-8. The neutrophils amplify the tissue inflammation. If stress and strain reach the limit of rupture of the fibre system, mechanical failure may

occur with direct rupture of alveolar walls and pulmonary capillaries. The excessive regional stress/strain may be limited by prone positioning and positive end-expiratory pressure (in a recruitable lung) which both allow a more uniform distribution of the stress and strain. A low global transpulmonary pressure decreases stress and strain and the available clinical evidence suggests that a low tidal volume is associated with less ventilator-induced lung injury than a high tidal volume. It would have been better, however, to test different plateau pressures, which are linked to the transpulmonary pressure more than to the tidal volume. The best solution, however, would be to directly test different transpulmonary pressures.

### References

1. Pontoppidan H, Geffin B, Lowenstein E. Acute respiratory failure in the adult. 3. *N Engl J Med* 1972; 287: 799–806.
2. Gattinoni L, Pesenti A, Avalli L, Rossi F, Bombino M. Pressure-volume curve of total respiratory system in acute respiratory failure. Computed tomographic scan study. *Am Rev Respir Dis* 1987; 136: 730–736.
3. Hickling KG, Henderson SJ, Jackson R. Low mortality associated with low volume pressure limited ventilation with permissive hypercapnia in severe adult respiratory distress syndrome. *Intensive Care Med* 1990; 16: 372–377.
4. Tremblay L, Valenza F, Ribeiro SP, Li J, Slutsky AS. Injurious ventilatory strategies increase cytokines and *c-fos* m-RNA expression in an isolated rat lung model. *J Clin Invest* 1997; 99: 944–952.
5. Ranieri VM, Suter PM, Tortorella C, *et al.* Effect of mechanical ventilation on inflammatory mediators in patients with acute respiratory distress syndrome: a randomized controlled trial. *JAMA* 1999; 282: 54–61.
6. Dreyfuss D, Ricard JD, Saumon G. On the physiologic and clinical relevance of lung-borne cytokines during ventilator-induced lung injury. *Am J Respir Crit Care Med* 2003; 167: 1467–1471.
7. Pelosi P, Cereda M, Foti G, Giacomini M, Pesenti A. Alterations of lung and chest wall mechanics in patients with acute lung injury: effects of positive end-expiratory pressure. *Am J Respir Crit Care Med* 1995; 152: 531–537.
8. Suter PM, Fairley HB, Isenberg MD. Effect of tidal volume and positive end-expiratory pressure on compliance during mechanical ventilation. *Chest* 1978; 73: 158–162.
9. Katz JA, Zinn SE, Ozanne GM, Fairley HB. Pulmonary, chest wall, and lung-thorax elastances in acute respiratory failure. *Chest* 1981; 80: 304–311.
10. Jardin F, Genevray B, Brun-Ney D, Bourdarias JP. Influence of lung and chest wall compliances on transmission of airway pressure to the pleural space in critically ill patients. *Chest* 1985; 88: 653–658.
11. Gattinoni L, Pelosi P, Suter PM, Pedoto A, Vercesi P, Lissoni A. Acute respiratory distress syndrome caused by pulmonary and extrapulmonary disease. Different syndromes? *Am J Respir Crit Care Med* 1998; 158: 3–11.
12. Ranieri VM, Brienza N, Santostasi S, *et al.* Impairment of lung and chest wall mechanics in patients with acute respiratory distress syndrome: role of abdominal distension. *Am J Respir Crit Care Med* 1997; 156: 1082–1091.
13. Weibel ER. Functional morphology of lung parenchyma. In: American Physiological Society, ed. *Handbook of Physiology: a Critical, Comprehensive Presentation of Physiological Knowledge and Concepts*. Baltimore, MD, Waverly press, Inc., 1986; pp. 89–111.
14. Wilson TA. Solid mechanics. In: American Physiological Society, ed. *Handbook of Physiology: a Critical, Comprehensive Presentation of Physiological Knowledge and Concepts*. Bethesda, MD, Waverly Press, Inc., 1986; pp. 35–39.
15. Bachofen H, Schurch S. Alveolar surface forces and lung

- architecture. *Comp Biochem Physiol A Mol Integr Physiol* 2001; 129: 183–193.
16. Mead J, Takishima T, Leith D. Stress distribution in lungs: a model of pulmonary elasticity. *J Appl Physiol* 1970; 28: 596–608.
  17. Muscedere JG, Mullen JB, Gan K, Slutsky AS. Tidal ventilation at low airway pressures can augment lung injury. *Am J Respir Crit Care Med* 1994; 149: 1327–1334.
  18. D'Angelo E, Pecchiari M, Baraggia P, Saetta M, Balestro E, Milic-Emili J. Low-volume ventilation causes peripheral airway injury and increased airway resistance in normal rabbits. *J Appl Physiol* 2002; 92: 949–956.
  19. Maksym GN, Bates JH. A distributed nonlinear model of lung tissue elasticity. *J Appl Physiol* 1997; 82: 32–41.
  20. Maksym GN, Fredberg JJ, Bates JH. Force heterogeneity in a two-dimensional network model of lung tissue elasticity. *J Appl Physiol* 1998; 85: 1223–1229.
  21. Weibel ER. *The Pathway for Oxygen Structure and Function in the Mammalian Respiratory System*. Cambridge, MA, Harvard University Press, 1984; pp. 303–338.
  22. Edwards YS. Stretch stimulation: its effects on alveolar type II cell function in the lung. *Comp Biochem Physiol A Mol Integr Physiol* 2001; 129: 245–260.
  23. Vlahakis NE, Hubmayr RD. Invited review: plasma membrane stress failure in alveolar epithelial cells. *J Appl Physiol* 2000; 89: 2490–2496.
  24. Vlahakis NE, Hubmayr RD. Response of alveolar cells to mechanical stress. *Curr Opin Crit Care* 2003; 9: 2–8.
  25. Liu M, Tanswell AK, Post M. Mechanical force-induced signal transduction in lung cells. *Am J Physiol* 1999; 277: L667–L683.
  26. Dos Santos CC, Slutsky AS. Invited review: mechanisms of ventilator-induced lung injury: a perspective. *J Appl Physiol* 2000; 89: 1645–1655.
  27. Uhlig S. Ventilation-induced lung injury and mechanotransduction: stretching it too far? *Am J Physiol Lung Cell Mol Physiol* 2002; 282: L892–L896.
  28. Pugin J. Molecular mechanisms of lung cell activation induced by cyclic stretch. *Crit Care Med* 2003; 31: S200–S206.
  29. Bachofen H, Schurch S, Urbinelli M, Weibel ER. Relations among alveolar surface tension, surface area, volume, and recoil pressure. *J Appl Physiol* 1987; 62: 1878–1887.
  30. Vlahakis NE, Schroeder MA, Pagano RE, Hubmayr RD. Role of deformation-induced lipid trafficking in the prevention of plasma membrane stress failure. *Am J Respir Crit Care Med* 2002; 166: 1282–1289.
  31. Vlahakis NE, Schroeder MA, Pagano RE, Hubmayr RD. Deformation-induced lipid trafficking in alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 2001; 280: L938–L946.
  32. Gajic O, Lee J, Doerr CH, Berrios JC, Myers JL, Hubmayr RD. Ventilator-induced cell wounding and repair in the intact lung. *Am J Respir Crit Care Med* 2003; 167: 1057–1063.
  33. Dudek SM, Garcia JG. Cytoskeletal regulation of pulmonary vascular permeability. *J Appl Physiol* 2001; 91: 1487–1500.
  34. Pugin J, Dunn I, Jolliet P, et al. Activation of human macrophages by mechanical ventilation *in vitro*. *Am J Physiol* 1998; 275: L1040–L1050.
  35. Haseneen NA, Vaday GG, Zucker S, Foda HD. Mechanical stretch induces MMP-2 release and activation in lung endothelium: role of EMMPRIN. *Am J Physiol Lung Cell Mol Physiol* 2003; 284: L541–L547.
  36. Yamamoto H, Teramoto H, Uetani K, Igawa K, Shimizu E. Cyclic stretch upregulates interleukin-8 and transforming growth factor- $\beta$ 1 production through a protein kinase C-dependent pathway in alveolar epithelial cells. *Respirology* 2002; 7: 103–109.
  37. Vlahakis NE, Schroeder MA, Limper AH, Hubmayr RD. Stretch induces cytokine release by alveolar epithelial cells *in vitro*. *Am J Physiol* 1999; 277: L167–L173.
  38. Tschumperlin DJ, Margulies SS. Equibiaxial deformation-induced injury of alveolar epithelial cells *in vitro*. *Am J Physiol* 1998; 275: L1173–L1183.
  39. Casetti AV, Bartlett RH, Hirschl RB. Increasing inspiratory time exacerbates ventilator-induced lung injury during high-pressure/high-volume mechanical ventilation. *Crit Care Med* 2002; 30: 2295–2299.
  40. Tschumperlin DJ, Oswari J, Margulies AS. Deformation-induced injury of alveolar epithelial cells. Effect of frequency, duration, and amplitude. *Am J Respir Crit Care Med* 2000; 162: 357–362.
  41. Hotchkiss JR Jr, Blanch L, Naveira A, et al. Relative roles of vascular and airspace pressures in ventilator-induced lung injury. *Crit Care Med* 2001; 29: 1593–1598.
  42. Mascheroni D, Kolobow T, Fumagalli R, Moretti MP, Chen V, Buckhold D. Acute respiratory failure following pharmacologically induced hyperventilation: an experimental animal study. *Intensive Care Med* 1988; 15: 8–14.
  43. Belperio JA, Keane MP, Burdick MD, et al. Critical role for CXCR2 and CXCR2 ligands during the pathogenesis of ventilator-induced lung injury. *J Clin Invest* 2002; 110: 1703–1716.
  44. Egan EA, Nelson RM, Olver RE. Lung inflation and alveolar permeability to non-electrolytes in the adult sheep *in vivo*. *J Physiol* 1976; 260: 409–424.
  45. Dreyfuss D, Basset G, Soler P, Saumon G. Intermittent positive-pressure hyperventilation with high inflation pressures produces pulmonary microvascular injury in rats. *Am Rev Respir Dis* 1985; 132: 880–884.
  46. Parker JC, Townsley MI, Rippe B, Taylor AE, Thigpen J. Increased microvascular permeability in dog lungs due to high peak airway pressures. *J Appl Physiol* 1984; 57: 1809–1816.
  47. Woo SW, Hedley-Whyte J. Macrophage accumulation and pulmonary edema due to thoracotomy and lung over inflation. *J Appl Physiol* 1972; 33: 14–21.
  48. Kolobow T, Moretti MP, Fumagalli R, et al. Severe impairment in lung function induced by high peak airway pressure during mechanical ventilation. An experimental study. *Am Rev Respir Dis* 1987; 135: 312–315.
  49. Webb HH, Tierney F. Experimental pulmonary edema due to intermittent positive pressure ventilation with high inflation pressures. Protection by positive end-expiratory pressure. *Am Rev Respir Dis* 1974; 110: 556–565.
  50. Parker JC, Hernandez LA, Longenecker GL, Peevy K, Johnson W. Lung edema caused by high peak inspiratory pressures in dogs. Role of increased microvascular filtration pressure and permeability. *Am Rev Respir Dis* 1990; 142: 321–328.
  51. Fu Z, Costello ML, Tsukimoto K, et al. High lung volume increases stress failure in pulmonary capillaries. *J Appl Physiol* 1992; 73: 123–133.
  52. Mathieu-Costello O, Willford DC, Fu Z, Garden RM, West JB. Pulmonary capillaries are more resistant to stress failure in dogs than in rabbits. *J Appl Physiol* 1995; 79: 908–917.
  53. West JB. Invited review: pulmonary capillary stress failure. *J Appl Physiol* 2000; 89: 2483–2489.
  54. Michel CC, Curry FE. Microvascular permeability. *Physiol Rev* 1999; 79: 703–761.
  55. Parker JC, Ivey CL, Tucker JA. Gadolinium prevents high airway pressure-induced permeability increases in isolated rat lungs. *J Appl Physiol* 1998; 84: 1113–1118.
  56. Parker JC. Inhibitors of myosin light chain kinase and phosphodiesterase reduce ventilator-induced lung injury. *J Appl Physiol* 2000; 89: 2241–2248.
  57. Parker JC, Hernandez LA, Peevy KJ. Mechanisms of ventilator-induced lung injury. *Crit Care Med* 1993; 21: 131–143.
  58. Broccard AF, Vannay C, Feihl F, Schaller MD. Impact of low pulmonary vascular pressure on ventilator-induced lung injury. *Crit Care Med* 2002; 30: 2183–2190.
  59. Broccard A, Shapiro RS, Schmitz LL, Adams AB, Nahum A, Marini JJ. Prone positioning attenuates and redistributes

- ventilator-induced lung injury in dogs. *Crit Care Med* 2000; 28: 295–303.
60. Nishimura M, Honda O, Tomiyama N, Johkoh T, Kagawa K, Nishida T. Body position does not influence the location of ventilator-induced lung injury. *Intensive Care Med* 2000; 26: 1664–1669.
  61. Guo F, Qiu H, Tan Y. [Low flow technique to perform static pressure-volume curve during mechanical ventilation.]. *Zhonghua Jie He He Hu Xi Za Zhi* 2001; 24: 728–731.
  62. Reta GS, Riva JA, Piriz H, Medeiros AS, Rocco PR, Zin WA. Effects of halothane on respiratory mechanics and lung histopathology in normal rats. *Br J Anaesth* 2000; 84: 372–377.
  63. Allen G, Lundblad LK, Parsons P, Bates JH. Transient mechanical benefits of a deep inflation in the injured mouse lung. *J Appl Physiol* 2002; 93: 1709–1715.
  64. Mercer RR, Russell ML, Crapo JD. Alveolar septal structure in different species. *J Appl Physiol* 1994; 77: 1060–1066.
  65. Dreyfuss D, Soler P, Basset G, Saumon G. High inflation pressure pulmonary edema. Respective effects of high airway pressure, high tidal volume, and positive end-expiratory pressure. *Am Rev Respir Dis* 1988; 137: 1159–1164.
  66. Dreyfuss D, Soler P, Saumon G. Mechanical ventilation-induced pulmonary edema. Interaction with previous lung alterations. *Am J Respir Crit Care Med* 1995; 151: 1568–1575.
  67. Hernandez LA, Coker PJ, May S, Thompson AL, Parker JC. Mechanical ventilation increases microvascular permeability in oleic acid-injured lungs. *J Appl Physiol* 1990; 69: 2057–2061.
  68. Gattinoni L, Pesenti A, Baglioni S, Vitale G, Rivolta M, Pelosi P. Inflammatory pulmonary edema and positive end-expiratory pressure: correlations between imaging and physiologic studies. *J Thorac Imaging* 1988; 3: 59–64.
  69. Gattinoni L, D'Andrea L, Pelosi P, Vitale G, Pesenti A, Fumagalli R. Regional effects and mechanism of positive end-expiratory pressure in early adult respiratory distress syndrome. *JAMA* 1993; 269: 2122–2127.
  70. Held HD, Boettcher S, Hamann L, Uhlig S. Ventilation-induced chemokine and cytokine release is associated with activation of nuclear factor- $\kappa$ B and is blocked by steroids. *Am J Respir Crit Care Med* 2001; 163: 711–716.
  71. Pelosi P, Goldner M, McKibben A, et al. Recruitment and derecruitment during acute respiratory failure: an experimental study. *Am J Respir Crit Care Med* 2001; 164: 122–130.
  72. Gattinoni L, Pelosi P, Valenza F, Mascheroni D. Patient positioning in acute respiratory failure. In: Tobin M ed. *Principles and Practice of Mechanical Ventilation*. New York, NY, McGraw-Hill, 1994; pp. 1067–1077.
  73. Mutoh T, Guest RJ, Lamm WJ, Albert RK. Prone position alters the effect of volume overload on regional pleural pressures and improves hypoxemia in pigs *in vivo*. *Am Rev Respir Dis* 1992; 146: 300–306.
  74. Gattinoni L, Tognoni G, Pesenti A, et al. Effect of prone positioning on the survival of patients with acute respiratory failure. *N Engl J Med* 2001; 345: 568–573.
  75. Dreyfuss D, Saumon G. Ventilator-induced lung injury: lessons from experimental studies. *Am J Respir Crit Care Med* 1998; 157: 294–323.
  76. Slutsky AS. Consensus conference on mechanical ventilation – January 28–30, 1993 at Northbrook, Illinois, USA. Part 2. *Intensive Care Med* 1994; 20: 150–162.
  77. Stewart TE, Meade MO, Cook DJ, et al. Evaluation of a ventilation strategy to prevent barotrauma in patients at high risk for acute respiratory distress syndrome. Pressure- and Volume-Limited Ventilation Strategy Group. The Multi-center Trial Group on Tidal Volume Reduction in ARDS. *N Engl J Med* 1998; 338: 355–361.
  78. Amato MB, Barbas CS, Medeiros DM, et al. Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. *N Engl J Med* 1998; 338: 347–354.
  79. Brochard L, Roudot-Thoraval F, Roupie E, et al. Tidal volume reduction for prevention of ventilator-induced lung injury in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1998; 158: 1831–1838.
  80. Brower RG, Shanholtz CB, Fessler HE, et al. Prospective, randomized, controlled clinical trial comparing traditional versus reduced tidal volume ventilation in acute respiratory distress syndrome patients. *Crit Care Med* 1999; 27: 1492–1498.
  81. The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000; 342: 1301–1308.
  82. Eichacker PQ, Gerstenberger EP, Banks SM, Cui X, Natanson C. Meta-analysis of acute lung injury and acute respiratory distress syndrome trials testing low tidal volumes. *Am J Respir Crit Care Med* 2002; 166: 1510–1514.
  83. Tobin MJ. Culmination of an era in research on the acute respiratory distress syndrome. *N Engl J Med* 2000; 342: 1360–1361.